

FOR ALL PROCEDURES

PRIOR TO STOPPING ANY MEDICATION, PLEASE GET APPROVAL FROM YOUR PRIMARY CARE PHYSICIAN.

1. Stop all **aspirin and aspirin containing medications, Plavix, Aggrenox** and other anti-platelet agents at least **5 days** prior to procedure.
2. Stop all anti-inflammatory medications such as **ibuprofen, naproxen, Aleve, Motrin**, etc., at least **48 hours** prior to procedures. Usually Celebrex, Bextra or Vioxx do not need to be stopped, but if you have any questions, please check with your doctor.
3. Stop **Vitamin E and Ginko Biloba** at least **5 days** prior to procedures.
4. Stop **Coumadin** at least **5 days** prior to procedures.
5. Stop low molecular weight heparin compounds such as **Lovenox** at least **24 hours** prior to procedure.
6. Stop all prescriptive or herbal **diet medication** at least **24 hours** prior to procedure.
7. If you are diabetic and/or take diabetes medication, please check with your doctor.

IF YOU ARE HAVING SEDATION FOR YOUR PRODECURE:

1. You must have an escort drive you home following your procedure. This person must be identified to the staff prior to the start of the procedure.
2. Cedars-Sinai policy prohibits our discharging you after sedation if you are unaccompanied this includes leaving our office via cab, bus or car service.
3. You may not have anything to eat for **8 hours** prior to your procedure. Nothing to drink including water, **6 hours** prior to the procedure.
4. If you usually take medication in the morning, please take it with just sips of water.